



The surveillance programme for small ruminant lentivirus infections in sheep and goats in Norway 2020



REPORT 17/2021

The surveillance programme for small ruminant lentivirus infections in sheep and goats in Norway 2020

Authors

Annette H Kampen, Johan Åkerstedt, Siv Klevar

Suggested citation

Kampen, Annette H, Åkerstedt, Johan, Klevar Siv. The surveillance programme for small ruminant lentivirus infections in sheep and goats in Norway 2020. Surveillance program report. Veterinærinstituttet 2021. © Norwegian Veterinary Institute, copy permitted with citation

Quality controlled by

Merete Hofshagen, Director of Animal Health, Animal Welfare and Food Safety, Norwegian Veterinary Institute

Published

2021 on www.vetinst.no
ISSN 1890-3290 (electronic edition)
© Norwegian Veterinary Institute 2021

Commissioned by / In collaboration with

Norwegian Food Safety Authority



Colophon

Cover design: Reine Linjer
Cover photo: Colourbox
www.vetinst.no

Content

Summary	3
Introduction	3
Aims	4
Materials and methods	4
Results	5
Sheep	5
Goat	6
Discussion	7
References.....	8

Summary

None of the 2,927 investigated sheep flocks in the surveillance programme were diagnosed with maedi in 2020. Of 50 tested goat herds, no herds were diagnosed with caprine arthritis-encephalitis (CAE).

Introduction

Small ruminant lentivirus (SRLV) comprise the maedi-visna virus (MVV) and the caprine arthritis-encephalitis virus (CAEV), which are closely related. Maedi-visna virus mainly infects sheep, while CAEV mainly infects goats, although cross-infections between the two species do occur. Maedi is a progressive viral pneumonia in sheep first described in Iceland in 1939 (1). The disease occurs in several European countries as well as in other continents. The disease visna is a neuropathogenic manifestation of the infection (1, 2). Caprine arthritis-encephalitis (CAE) causes emaciation, arthritis, encephalitis and sometimes mastitis and pneumonia in goats. Sheep may be infected and produce antibodies against CAEV, but usually show no clinical signs of disease.

Maedi-visna and CAE are classified as list B diseases in Norway and are notifiable to the Office International des Epizooties. In Norway, maedi was officially reported for the first time in 1972 (3).

In November 2002 and January 2003, post-mortem examinations of lungs from two diseased sheep from two different farms in the former Nord-Trøndelag county showed histopathological changes consistent with maedi. During the following investigations more than 15,000 sheep in 300 flocks were serologically examined for maedi-visna infection, and approximately 50 flocks were found to be seropositive (4, 5). The outbreak demonstrated the need for a nationwide surveillance programme, and up to 2018 no new cases of maedi were detected (4, 6)

In 2019, the surveillance programme detected a new case of maedi in a farm in Trøndelag county. Contacts to this farm were traced, and an outbreak zone was established in Northern Trøndelag, where all sheep farms were monitored by serology. A total of nine herds has so far been diagnosed with maedi in the outbreak (G. Rømo, manuscript in preparation).

An overview of the number of new infected flocks registered with maedi each year up to 2020 is given in Figure 1.




Figure 1: The number of new flocks infected with maedi from 1972 and onwards. The bars for 2003 - 2005 show both seropositive flocks detected through the investigations after the outbreak in the former Nord-Trøndelag county and seropositive flocks identified in the programme. The rise in 2019 represents a new maedi outbreak in Trøndelag.

Before 2001, CAE was widespread in the Norwegian dairy goat population. The dairy organisation (TINE) and the Norwegian Goat Health Services have conducted an eradication programme named “Healthier goats”, targeting caprine arthritis encephalitis, caseous lymphadenitis and paratuberculosis. The programme started in 2001, and in total 612 goat herds were included in the programme from 2001 to 2014 (7). In 2018, the goat population was included in the surveillance programme for SRLV.

Aims

The aims of the surveillance programme for small ruminant lentivirus infections are to document the status for NVV and CAEV infections in sheep and goats in Norway and to identify infected flocks for disease control.

Materials and methods

The surveillance programme is based on serological examination for SRLV in both sheep and goats.

The plan for 2020, was to collect 9 000 blood samples from sheep at slaughter. Representative numbers of animals were sampled at 22 larger abattoirs, from various parts of Norway. A maximum of five animals (>2 years old) were to be sampled per herd any given day. The preferred sampling period was from January to May, but a proportion of the animals were sampled from September to November. In addition, meat inspectors at the abattoirs were asked to monitor sheep and especially their lungs for detection of suspicious cases consistent with maedi.

In addition, 60 goat herds were randomly selected for sampling. In goat herds of less than 30 animals, all animals (>2 years old) were sampled. In herds of 30 to 100, 100 to 200, and more than 200 animals, samples from 30, 35, and 40 animals were sampled, respectively.

Serum samples were examined for antibodies against SRLV with ID Screen® MVV / CAEV Indirect ELISA (IDvet, Grabels, France) for initial screening. Samples with inconclusive or positive ELISA results were re-tested in duplicates with ID Screen® MVV / CAEV Indirect ELISA (IDvet). If samples were inconclusive or positive, samples were further tested in duplicates with ELISA IDEXX MVV/CAEV p28 Ab Verification Test (IDEXX Laboratories, Maine, USA).

In case of positive or inconclusive results in IDEXX ELISA on a sample taken from a sheep at slaughter, follow up sampling was done on selected animals in the flock of origin as described previously (8).

Samples with doubtful or positive status in confirmatory tests were reported, and new blood samples from the suspected animals or herd were requested and tested.

Results

Sheep

A total of 8,701 samples from 2,927 sheep flocks were received in the programme in 2020 (Table 1). This was approximately 21.7% of the total number of Norwegian sheep flocks.

In 2020 all sheep samples were concluded to be negative for antibodies against SRLV. A total of 49 samples from 46 herds tested positive with the IDvet assay, whereas the IDEXX ELISA gave negative results for all these samples.

Sixteen farms in five counties with positive results in the screening assay, were followed-up by sampling according to the sampling protocol given for goats in the Materials and Methods section. All 528 additionally sampled animals from these 16 farms tested negative. Thus, maedi was not suspected in these farms.

Lung samples from five suspected cases of maedi from four farms were submitted by meat inspectors at abattoirs. Pathological examination and PCR for SRLV were negative. In each herd of origin, 30 - 50 animals were sampled and found to be serologically negative.

Tabell 1: The results and total number of sheep flocks within the frame of the Norwegian surveillance programme for maedi 2003-2020.

Year	Total no. of flocks*	No. of flocks analysed	No. of animals analysed	Average no. of animals analysed per flock	No. of positive flocks
2003	18 400	456**	13 951	30.6	1
2004	17 439	1 230	36 911	30.0	1
2005	16 500	940	29 248	31.1	2
2006	15 800	911	27 846	30.6	0
2007	15 400	1 004	29 633	29.5	0
2008	15 059	783	23 235	29.7	0
2009	14 800	417	12 198	29.3	0
2010	14 800	188	5 697	60.6	0
2011	14 500	467	13 628	29.2	0
2012	14 300	479	14 043	29.3	0
2013	14 242	468	13 550	29.0	0
2014	14 218	3 506	9 771	2.8	0
2015	14 425	3 357	9 442	2.8	0
2016	14 561	3 504	9 858	2.8	0
2017	14 463	3 447	9 041	2.6	0
2018	14 337	3 282	8 685	2.6	0
2019	13 693	3 264	8 993	2.6	1
2020	13 507	2 927	8 701	3.0	0

* Based on data from the register of production subsidies as of 31st of July the respective year until 2017. Thereafter, data from 1st of March was used.

** Sampling period: 20th of November to 31st of December.

Goats

A total of 1,443 samples from 50 goat herds were received and tested for antibodies against CAEV (Table 2). This was approximately 3.9% of Norwegian goat herds.

All 1,443 samples tested negative for SRLV.

Tabell 2: The results and total number of goat flocks within the frame of the Norwegian surveillance programme for small ruminant lentivirus 2020.

Year	Total no. of flocks*	No. of flocks analysed	No. of animals analysed	Average no. of animals analysed per flock	No. of positive flocks
2018	1 246	61	1 663	27.2	1
2019	1 209	58	1 751	30.2	1
2020	1 271	50	1 443	28.9	0

* Based on data from the register of production subsidies as of 1st of March the respective year.

Suspicion of CAE was raised two other goat herds in Møre og Romsdal county based on milk sample testing at the dairy organisation (TINE) laboratory in Molde, which is not part of the surveillance programme run by the food safety authorities. In the two herds, 54 and 189 animals (>1 year old) were sampled, respectively. One sample in each herd was positive for antibodies against SRLV. The two flocks were concluded to be infected with CAEV.

Discussion

During the years 2003-2008, ram circles and their flocks registered as members of The Norwegian Association of Sheep and Goat Farmers constituted the target population for the programme. Approximately 90% of the Norwegian sheep flocks participating in ram circles were screened for antibodies against MVV during 2003 to 2005. These herds were retested during 2006 to 2008. In 2009, breeding flocks of other sheep breeds than those represented by The Norwegian Sheep and Goat Breeders Association were selected for sampling. During 2010-2013, randomly selected sheep flocks were sampled.

From 2014, sheep have been sampled at slaughterhouses, giving a better surveillance of the total population with the use of less resources than needed when sampling on farms (8). However, less animals were tested per herd, giving less accurate results on the herd level.

In goats, the surveillance is still based on sampling live animals in the herds.


Results from the surveillance programme for maedi from November 2003 through 2006, showed a prevalence of less than 0.2% positive flocks. The distribution of the disease indicated that it was regionally clustered. A more extensive spread of MVV virus from the outbreak in 2003 was probably prevented by the restrictions on transfer of sheep across county borders. Maedi was not detected in the surveillance programme in the years 2005 to 2018, which indicated that the prevalence of the infection in Norway was very low. Even if maedi has been detected in nine farms in total during 2019 and 2020, the prevalence of infected herds still remains very low (less than 0.07%).

The design of the surveillance programme with testing mainly flocks in the breeding system the first six years, made it possible for maedi infection to remain undetected in other types of flocks. The sampling design since 2014 of few samples from a large number of flocks resulted in the detection of one flock in the surveillance programme in 2019. Another eight infected flocks were found by tracing of contacts and extended screening, some of which had probably been infected for years. To achieve higher sensitivity of the programme, more animals and more flocks would need to be tested. Also, sampling could be targeted on clinical signs consistent with maedi infection, and to geographical areas in Norway where maedi has occurred in the past.

All dairy goat herds in Norway and many meat and fiber goat herds have joined the eradication programme Healthier goats. This has resulted in improved goat health and welfare in the Norwegian goat industry (7, 9). The finding of CAE in several hobby goat herds in 2019 as well as in two dairy goat herds in 2020 where CAE previously had been eradicated, shows that attention and continued surveillance is needed in all types of goat herds.

References

1. Pálsson PA. Maedi-visna. History and clinical description. In: Pétursson G, Hoff-Jørgensen R (editors). *Maedi-visna and Related Diseases*. Boston: Kluwer Academic Publishers; 1990. p. 3-17.
2. Martin WB, Aitken ID. *Diseases of Sheep*, 3rd edition. Oxford: Blackwell Scientific Publications; 2000.
3. Krogsrud J, Larsen HJS, Rimstad E. Mædi og lungeadenomatose [Maedi and lung adenomatosis, No]. *Nor Vet Tidsskr.* 1996; 108: 729-36.
4. Sviland S, Nyberg O, Tharaldsen J, Heier B T, Mork J. The surveillance and control programme for maedi in Norway. In: Mørk T, Hellberg H (editors). *Surveillance and control programmes for terrestrial and aquatic animals in Norway. Annual report 2003*. Oslo: National Veterinary Institute; 2004. p. 89-95.
5. Kampen AH, Tharaldsen J, Åkerstedt J, Norström M, Nestvold OK, Myhre JL, Nyberg O. Diagnosis and investigations of an outbreak of maedi in Norway 2002 - 2005. *Proceedings of the 6th International Veterinary Vaccines and Diagnostics Conference; Jun 25 - 29. Oslo, Norway; 2006*. p. 91-2.
6. Mork J, Jarp J. The surveillance and control programme for maedi in Norway. In: Fredriksen B, Mørk T (editors). *Surveillance and control programmes for terrestrial and aquatic animals in Norway. Annual report 2001*. Oslo: National Veterinary Institute; 2002. p. 109-15.
7. TINE Rådgiving, Helsetjenesten for geit. Syk - friskere - friskest. Sluttrapport prosjekt Friskere geiter 2001-2015. [Healthier goats project, final report 2001-2015, No]. Ås: Helsetjenesten for Geit; 2016.
8. Kampen AH, Åkerstedt J, Rømo G, Mork J, Nordstoga A, Klevar S. The surveillance programme for small ruminant lentivirus infections in sheep and goats in Norway 2019. *Surveillance programmes for terrestrial and aquatic animals in Norway. Annual report 2019*. Oslo: Norwegian Veterinary Institute; 2020.
9. Muri K, Leine N, Valle PS. Welfare effects of a disease eradication programme for dairy goats. *Animal.* 2016; 10: 333-41.



*Scientifically ambitious, forward-looking
and collaborative- for one health!*



Veterinærinstituttet
Norwegian Veterinary Institute

Ås

Trondheim

Sandnes

Bergen

Harstad

Tromsø

postmottak@vetinst.no
www.vetinst.no